arXiv:0904.4639v1 [hep-th] 29 Apr 2009

Quark-antiquark and diquark condensates in vacuum in two-flavor four-fermion interaction models with any color number N_c *

ZHOU Bang-Rong†

College of Physical Sciences, Graduate University of the
Chinese Academy of Sciences, Beijing 100049, China

(Dated:)

The color number N_c-dependence of the interplay between quark-antiquark condensates $\langle\bar{q}q\rangle$ and diquark condensates $\langle qq\rangle$ in vacuum in two-flavor four-fermion interaction models is researched. The results show that the G_S-H_S (the coupling constant of scalar $(\bar{q}q)^2$-scalar $(qq)^2$ channel) phase diagrams will be qualitatively consistent with the case of $N_c = 3$ as N_c varies in 4D Nambu-Jona Lasinio model and 2D Gross-Neveu (GN) model. However, in 3D GN model, the behavior of the G_S-H_P (the coupling constant of pseudoscalar $(qq)^2$ channel) phase diagram will obviously depend on N_c. The known characteristic that a 3D GN model has not the coexistence phase of the condensates $\langle\bar{q}q\rangle$ and $\langle qq\rangle$ is proven to appear only in the case of $N_c \leq 4$. In all the models, the regions occupied by the phases containing the diquark condensates $\langle qq\rangle$ in corresponding phase diagrams will gradually decrease as N_c grows up and finally go to zero if $N_c \to \infty$, i.e. in this limit only the pure $\langle\bar{q}q\rangle$ phase could exist.

PACS numbers: 11.10.Lm, 11.10.Wx, 11.15.Pg, 11.30.Rd

Keywords: four-fermion interaction model, quark-antiquark and diquark condensates, color number N_c

I. INTRODUCTION

The four-fermion interaction models including 4D Nambu-Jona-Lasinio (NJL) model [1], 3D and 2D Gross-Neveu (GN) model [2] are not only ideal theoretical laboratories to research dynamical symmetry breaking [1, 2, 3, 4, 5, 6] and in recent years, the color-

---

[*] The project supported by the National Natural Science Foundation of China under Grant No.10475113.

[†] E-mail:zhoubr@gucas.ac.cn

superconductivity [7, 8], but also can find many practical usefulness to Particle Physics and Condensed Matter Physics. Hence it is certainly interesting to conduct some more deepgoing exploration for these field theory models themselves.

Let q denote the fermion (called as quark) field and $\bar{q}$ antiquark field. Since in any four-fermion interaction model, the couplings of $(qq)^2$-form and $(\bar{q}q)^2$-form can always coexist via the Fierz transformation [8], so once a sufficiently strong coupling of $(qq)^2$-form leads to formation of the diquark condensates $\langle qq\rangle$, then there will certainly exist interplay between the quark-antiquark condensates $\langle\bar{q}q\rangle$ and the diquark condensates $\langle qq\rangle$ in the ground state of the system, even in vacuum [9, 10, 11]. In vacuum condition, despite of absence of net quarks, however, based on the relativistic quantum field theory, it is possible that the diquark condensates $\langle qq\rangle$ and di-antiquark condensates $\langle\bar{q}\bar{q}\rangle$ are generated simultaneously.

Motivated on the above idea, we have researched interplay between the condensates $\langle\bar{q}q\rangle$ and $\langle qq\rangle$ in vacuum in two-flavor 4D NJL [12], 2D GN [13] and 3D GN [14] models and find that the ratio between the coupling constants of the $(\bar{q}q)^2$-form and the $(qq)^2$-form plays a key role in determining the property of the ground state. It should be indicated that some properties of the system in the vacuum could be maintained to the case of finite temperature and density. For instance, it has been shown that [15], in a 3D GN model with the quark's color number $N_c = 3$, the conclusion derived from the vacuum condition that the model does not contain the phase with coexistence of the condensates $\langle\bar{q}q\rangle$ and $\langle qq\rangle$ will be kept to the case of finite temperature and finite quark chemical potential. Therefore, the property of a model in vacuum could reflect its some essential characteristic and this fact will make the research on interplay between the two condensates in vacuum be given more theoretical significance. In addition, a technical advantage is that all the discussions can be made in an analytical way in this case.

In our preceding work, the color number N_c is fixed to be 3, consistent with phenomenology of Quantum Chromodynamics (QCD). In this paper, we will extend the research to more general case of any N_c. In fact, the large N_c behavior of a four-fermion interaction model is always an interesting problem to deserve to be followed.

We will still use the effective potential approach employed in Refs.[12, 13, 14] and obtain the expressions of relevant effective potentials by similar procedures but omit the details of these procedures. The mean field approximation will be taken. In this case we will use the Fierz transformed four-fermion couplings in the Hartree approximation so as to avoid double counting [8]. In selecting the couplings of the $(qq)^2$-form, we will always simulate the $SU_c(N_c)$ gauge interactions, where two fermions are attractive each other in the antisymmetric $N_c(N_c-1)/2$-plet. The main results will also be shown in some figures more clearly.

II. 4D NAMBU-JONA-LASINIO MODEL WITH ANY N_c

With 2 flavor and N_c color massless fermions, the Lagrangian

$$\begin{aligned}\mathcal{L} &= \bar{q}i\gamma^\mu\partial_\mu q + G_S[(\bar{q}q)^2 + (\bar{q}i\gamma_5\vec{\tau}q)^2] \\ &\quad + H_S\sum_{\lambda_A}(\bar{q}i\gamma_5\tau_2\lambda_A q^C)(\bar{q}^C i\gamma_5\tau_2\lambda_A q), \end{aligned} \tag{1}$$

where the fermion fields q are in the doublet of $SU_f(2)$ and the N_c-plet of $SU_c(N_c)$, i.e.

$$q = \begin{pmatrix} u_i \\ d_i \end{pmatrix} \quad i = 1, \cdots, N_c, \tag{2}$$

q^C is the charge conjugate of q and $\vec{\tau} = (\tau_1, \tau_2, \tau_3)$ are the Pauli matrices acting in two-flavor space. The matrices λ_A run over all the antisymmetric generators of $SU_c(N_c)$, G_S and H_S are respectively the coupling constants in scalar $(\bar{q}q)^2$ channel and scalar color $N_c(N_c-1)/2$-plet $(qq)^2$ channel.

Assume that the four-fermion interactions can lead to the scalar condensates

$$\langle\bar{q}q\rangle = \phi \tag{3}$$

with all the N_c color fermion entering them, and the scalar color $N_c(N_c-1)/2$-plet difermion and di-antifermion condensates (after a global $SU_c(N_c)$ transformation)

$$\langle\bar{q}^C i\gamma_5\tau_2\lambda_2 q\rangle = \delta, \quad \langle\bar{q}i\gamma_5\tau_2\lambda_2 q^C\rangle = \delta^*, \tag{4}$$

with only two color fermions enter them. The corresponding symmetry breaking is that $SU_{fL}(2)\otimes SU_{fR}(2) \to SU_f(2)$, $SU_c(N_c) \to SU_c(2)$, and a "rotated" electric charge $U_{\tilde{Q}}(1)$ and a "rotated" quark number $U'_q(1)$ leave unbroken [8]. It should be indicated that in the case of vacuum, the Goldstone bosons induced by spontaneous breaking of $SU_c(N_c)$ could be some combinations of difermions and di-antifermions [12].
Define that

$$\sigma = -2G_S\phi, \;\; \Delta = -2H_S\delta, \;\; \Delta^* = -2H_S\delta^*. \tag{5}$$

With a 4D Euclidean momentum cutoff Λ, we can obtain the relativistic effective potential [12]

$$\begin{aligned}V_4(\sigma,|\Delta|) &= \frac{\sigma^2}{4G_S} + \frac{|\Delta|^2}{4H_S} - \frac{1}{4\pi^2}\left[(N_c\sigma^2 + 2|\Delta|^2)\Lambda^2 - (N_c-2)\frac{\sigma^4}{2}\left(\ln\frac{\Lambda^2}{\sigma^2} + \frac{1}{2}\right)\right. \\ &\quad \left. -(\sigma^2+|\Delta|^2)^2\left(\ln\frac{\Lambda^2}{\sigma^2+|\Delta|^2} + \frac{1}{2}\right)\right], \end{aligned} \tag{6}$$



Fig.1: G_S-H_S phase diagram of a 4D Nambu-Jona-Lasinio model with the denotations that $(0, \Delta_1)$–pure $\langle qq\rangle$ phase, (σ_2, Δ_2)–mixed phase with $\langle \bar{q}q\rangle$ and $\langle qq\rangle$ and $(\sigma_1, 0)$–pure $\langle \bar{q}q\rangle$ phase.

where the conditions $\Lambda^2 \gg \sigma^2$ and $\Lambda^2 \gg \sigma^2+\Delta^2$ have been assumed. Through an analytic calculations, we have found that the ground states of the system, i.e. the minimum points of $V_4(\sigma, |\Delta|)$, will be at

$$(\sigma, |\Delta|) = \left\{ \begin{array}{ll} (0, & \Delta_1) \\ (\sigma_2, & \Delta_2) \\ (\sigma_1, & 0) \end{array} \right. \text{if} \left\{ \begin{array}{rcl} H_S\Lambda^2/\pi^2 > 1/2, \qquad 0 & \leq & G_S/H_S < 1/[1+(N_c-2)H_S\Lambda^2/\pi^2] \\ 1/[1+(N_c-2)H_S\Lambda^2/\pi^2] & < & G_S/H_S < 2/N_c \\ G_S\Lambda^2/\pi^2 > 1/N_c, & & G_S/H_S > 2/N_c \end{array} \right. , \tag{7}$$

Eq.(7) gives the phase diagram Fig.1 of the 4D NJL model.

The results qualitatively coincides with the ones obtained in Ref.[12] except that the color number is changed from 3 to N_c. Now the pure $\langle \bar{q}q\rangle$ phase happens only if the ratio $G_S/H_S > 2/N_c$. This conclusion was obtained in Ref.[5] but no $G_S - H_S$ phase diagram was given there. The $2/N_c$ is just the ratio of the color numbers of the quarks entering the diquark condensates and the quark-antiquark condensates. The diquark condensates $\langle qq\rangle$ could emergy only in the region where $H_S\Lambda^2/\pi^2 > 1/2$ and $G_S/H_S < 2/N_c$. It is seen from Fig.1 that as G_S/H_S decreases (or H_S increases), we first come to a phase with coexistence of the quark-antiquark condensates and the diquark condensates, then to a pure diquark condensate phase, the boundary between the two phases is determined by the curve $G_S = H_S/[1+(N_c-2)H_S\Lambda^2/\pi^2]$. On the other hand, for a fixed G_S and H_S, as N_c grows up, the slopes of the lines $G_S = 2H_S/N_c$ and $G_S = H_S/[1+(N_c-2)H_S\Lambda^2/\pi^2]$ will decrease. This implies that the regions with the diquark condensates will become smaller and smaller, and finally reduce to zeroes if $N_c \to \infty$. In the latter limit, only pure quark-antiquark condensates could exist.

III. 2D GROSS-NEVEU MODEL WITH ANY N_c

The Lagrangian of the model is expressed by

$$\begin{aligned} \mathcal{L} \;=\; & \bar{q}i\gamma^\mu\partial_\mu q + G_S[(\bar{q}q)^2 + (\bar{q}i\gamma_5\vec{\tau}q)^2] \\ & + H_S(\bar{q}i\gamma_5\tau_S\lambda_A q^C)(\bar{q}^C i\gamma_5\tau_S\lambda_A q), \end{aligned} \tag{8}$$

All the denotations are the same as ones in 4D NJL model, except that in 2D space-time

$$\gamma^0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \gamma^1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = -C, \quad \gamma_5 = \gamma^0\gamma^1$$

and $\tau_S = (\tau_0 \equiv 1, \tau_1, \tau_3)$ are flavor-triplet symmetric matrices. It is indicated that the product matrix $C\gamma_5\tau_S\lambda_A$ is antisymmetric.

Assume that the four-fermion interactions could lead to the scalar quark-antiquark condensates

$$\langle\bar{q}q\rangle = \phi, \tag{9}$$

which will break the discrete symmetries

$$\chi_D : q(t,x) \overset{\chi_D}{\to} \gamma_5 q(t,x),$$

$$\mathcal{P}_1 : q(t,x) \overset{\mathcal{P}_1}{\to} \gamma^1 q(t,-x),$$

and that the coupling with H_S can lead to the scalar color $N_c(N_c-1)/2$-plet difermion condensates and the scalar color anti- $N_c(N_c-1)/2$-plet di-antifermion condensates (after a global transformation in flavor and color space)

$$\langle\bar{q}^C i\gamma_5 1_f\lambda_2 q\rangle = \delta, \quad \langle\bar{q}i\gamma_5 1_f\lambda_2 q^C\rangle = \delta^* \tag{10}$$

which will break discrete symmetries $Z^c_{N_c}$ (center of $SU_c(N_c)$) down to Z^c_2, besides χ_D and $\mathcal{P}_1$[13]. Noting that in a 2D model, no breaking of continuous symmetry needs to be considered on the basis of Mermin-Wagner-Coleman theorem[16].

The model is renormalizable. We may derive the renormalized effective potential[13] containing the condensates (9) and (10). In the space-time dimension regularization approach, we can write down the renormalized $\mathcal{L}$ in $D = 2 - 2\varepsilon$ dimension space-time by the replacements

$$G_S \to G_S M^{2-D} Z_G, \; H_S \to H_S M^{2-D} Z_H,$$

with the scale parameter M, the renormalization constants Z_G and Z_H. In addition, the γ^μ in $\mathcal{L}$ will become $2^{D/2} \times 2^{D/2}$ matrices.

Define the order parameters

$$\sigma = -2G_S M^{2-D} Z_G \phi, \quad \Delta = -2H_S M^{2-D} Z_H \delta, \tag{11}$$

which will be finite if Z_G and Z_H are selected so as to cancel the UV divergences in ϕ and δ. In the minimal substraction scheme,

$$Z_G = 1 - \frac{2N_c G_S}{\pi}\frac{1}{\varepsilon}, \; Z_H = 1 - \frac{4H_S}{\pi}\frac{1}{\varepsilon}. \tag{12}$$

The corresponding renormalized effective potential in the mean field approximation up to one-loop order becomes

$$\begin{aligned} V_2(\sigma, |\Delta|) \; = \; & \frac{\sigma^2}{4G_S} - \frac{\sigma^2}{2\pi}\left(2\ln\frac{\bar{M}^2}{\sigma^2+|\Delta|^2} + (N_c-2)\ln\frac{\bar{M}^2}{\sigma^2} + N_c\right) \\ & + \frac{|\Delta|^2}{4H_S} - \frac{|\Delta|^2}{\pi}\left(\ln\frac{\bar{M}^2}{\sigma^2+|\Delta|^2} + 1\right), \; \bar{M}^2 = 2\pi e^{-\gamma} M^2, \end{aligned} \tag{13}$$

where γ is the Euler constant. $V_2(\sigma, |\Delta|)$ contains the two order parameters σ and $|\Delta|$. By almost identical demonstration to the one made in Ref.[13] (the only change is that the color number of quark varies from 3 to N_c), we obtain that the ground states of the system i.e. the minimal points of $V_2(\sigma, |\Delta|)$ will be at

$$(\sigma, |\Delta|) = \begin{cases} (0, \;\; \Delta_1) \\ (\sigma_2, \; \Delta_2) \\ (\sigma_1, \;\; 0) \end{cases} \text{if} \; \begin{cases} G_S/H_S = 0 \\ 0 < G_S/H_S < 2/N_c \\ G_S/H_S > 2/N_c \end{cases} \tag{14}$$

The results (14) can be drawn in Fig.2 which is the $G_S - H_S$ phase diagram of the 2D GN model.

The phase diagram are similar to the one obtained in the case of $N_c = 3$, including that formations of the condensates do not call for the coupling constant G_S and H_S having some lower bounds, however, the boundary between the pure quark-antiquark condensate phase and the coexistence phase with quark-antiquark condensate and diquark condensates is now replaced by $G_S/H_S = 2/N_c$. It is seen that as N_c increases, the area in Fig.2 occupied by the coexistence phase of quark-antiquark condensate and diquark condensates will gradually decrease and finally disappear if $N_c \to \infty$. It is indicated that if $N_c \to \infty$, the region corresponding to the pure diquark condensates in the diagram (the transverse axis of Fig.2) will also not exist. In fact, the pure diquark condensate phase corresponds to the minimal point of $V_2(\sigma, |\Delta|)$ $(\sigma, |\Delta|) = (0, \Delta_1)$, where Δ_1 is the



Fig.2: $G_S - H_S$ phase diagram of a 2D Gross-Neveu model with the denotations that $(0, \Delta_1)$ – pure $\langle qq \rangle$ phase, (σ_2, Δ_2) – mixed phase with $\langle \bar{q}q \rangle$ and $\langle qq \rangle$ and $(\sigma_1, 0)$ – pure $\langle \bar{q}q \rangle$ phase.

non-zero solution of the equation $\partial V_2(\sigma, |\Delta|)/\partial|\Delta| = 0$ with $\sigma = 0$. In this case, it is easy to find that the determinant of the second derivatives of V_2

$$K_2 = \begin{vmatrix} A_2 & B_2 \\ B_2 & C_2 \end{vmatrix}$$

with $A_2 = \partial^2 V_2/\partial\sigma^2$, $B_2 = \partial^2 V_2/\partial\sigma/\partial|\Delta| = \partial^2 V_2/\partial|\Delta|/\partial\sigma$ and $C_2 = \partial^2 V_2/\partial|\Delta|^2$. For the solution $(0, \Delta_1)$, we get that

$$A_2 = \frac{1}{2G_S} - \frac{1}{2H_S} - \frac{N_c - 2}{\pi} \ln \frac{\bar{M}^2}{\sigma^2}\bigg|_{\sigma\to 0} + \frac{2(N_c - 2)}{\pi},$$

$$K_2 = 4A_2/\pi.$$

If N_c is finite, then $G_S = 0$ could make $A_2 > 0$ and $K_2 > 0$ thus the solution $(0, \Delta_1)$ becomes a minimal point of V_2. However, if $N_c \to \infty$, then the third term of A_2 will have the $-\infty^2$ behavior, hence even if $G_S = 0$ then it is impossible to have $A_2 > 0$. As a result, the solution $(0, \Delta_1)$ is no longer a extreme point of V_2. This will imply that the region of the pure diquark condensate phase corresponding to the transverse axis of Fig.2 will not exist for finite H_S, $G_S = 0$ and $N_c \to \infty$.

IV. 3D GROSS-NEVEU MODEL WITH ANY N_c

The Lagrangian of the model is expressed by

$$\begin{aligned}\mathcal{L} &= \bar{q}i\gamma^\mu\partial_\mu q + G_S[(\bar{q}q)^2 + (\bar{q}\vec{\tau}q)^2] \\ &\quad + H_P\sum_{\lambda_A}(\bar{q}\tau_2\lambda_A q^C)(\bar{q}^C\tau_2\lambda_A q),\end{aligned} \tag{15}$$

where $\gamma^\mu(\mu = 0,1,2)$ are taken to be 2×2 matrices

$$\gamma^0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},\ \gamma^1 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix},\ \gamma^2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = C.$$

It is noted that the product matrix $C\tau_2\lambda_A$ is antisymmetric, and since without the "γ_5" matrix, the only possible color $N_c(N_c-1)/2$-plet difermion interaction channel is pseudoscalar one. The condensates $\langle\bar{q}q\rangle$ will break time reversal symmetry $\mathcal{T} : q(t,\vec{x}) \to \gamma^2 q(-t,\vec{x})$, special parity $\mathcal{P}_1 : q(t,x^1,x^2) \to \gamma^1 q(t,-x^1,x^2)$, special parity $\mathcal{P}_2 : q(t,x^1,x^2) \to \gamma^2 q(t,x^1,-x^2)$. The difermion condensates $\langle\bar{q}^C\tau_2\lambda_2 q\rangle$ (after a global rotation in the color space) will break $SU_c(N_c) \to SU_c(2)$ and leave a "rotated" electrical charge $U_{\tilde{Q}}(1)$ and a "rotated" fermion number $U'_q(1)$ unbroken. It also breaks parity $\mathcal{P} : q(t,\vec{x}) \to \gamma^0 q(t,-\vec{x})$ and this shows pseudoscalar feature of the difermion condensates.

Define the order parameters in the 3D GN model

$$\sigma = -2G_S\langle\bar{q}q\rangle, \quad \Delta = -2H_P\langle\bar{q}^C\tau_2\lambda_2 q\rangle, \tag{16}$$

The effective potential in the mean field approximation can be expressed by [14]

$$\begin{aligned}V_3(\sigma,|\Delta|) &= \frac{\sigma^2}{4G_S} + \frac{|\Delta|^2}{4H_P} - \frac{1}{\pi^2}(N_c\sigma^2 + 2|\Delta|^2)\Lambda_3 \\ &\quad + \frac{1}{3\pi}\left[6\sigma^2|\Delta| + 2|\Delta|^3 + (N_c-2)\sigma^3\right. \\ &\quad \left. + 2\theta(\sigma-|\Delta|)(\sigma-|\Delta|)^3\right],\end{aligned} \tag{17}$$

where Λ_3 is a 3D Euclidean momentum cutoff and the assumptions $\Lambda_3 \gg |\sigma - |\Delta||$, $\Lambda_3 \gg \sigma + |\Delta|$ and $\Lambda_3 \gg \sigma$ have been made. The ground states of the system correspond to the least value points of $V_3(\sigma,|\Delta|)$. Since the phase structure of the 3D GN model will have qualitatively different N_c-dependence from the one of the 4D and 2D models, we will make more detailed discussions of this problem.

The extreme value conditions $\partial V_3(\sigma,|\Delta|)/\partial\sigma = 0$ and $\partial V_3(\sigma,|\Delta|)/\partial|\Delta| = 0$ will respectively become

$$\begin{aligned}&\sigma\left(\frac{1}{2G_S} - \frac{2N_c\Lambda_3}{\pi^2} + \frac{4|\Delta|}{\pi} + \frac{(N_c-2)\sigma}{\pi}\right) \\ &+ \frac{2}{\pi}\theta(\sigma-|\Delta|)(\sigma-|\Delta|)^2 = 0,\end{aligned} \tag{18}$$

$$|\Delta|\left(\frac{1}{2H_P}-\frac{4\Lambda_3}{\pi^2}+\frac{2|\Delta|}{\pi}\right)$$
$$+\frac{2}{\pi}[\sigma^2-\theta(\sigma-|\Delta|)(\sigma-|\Delta|)^2]=0. \tag{19}$$

Define the determinant from the second derivatives of $V_3(\sigma,|\Delta|)$

$$K=\begin{vmatrix} A & B \\ B & C \end{vmatrix}=AC-B^2,$$

where

$$\begin{aligned} A &= \frac{\partial^2 V_3}{\partial\sigma^2}=\frac{1}{2G_S}-\frac{2N_c\Lambda_3}{\pi^2}+\frac{4|\Delta|}{\pi}+\frac{2(N_c-2)\sigma}{\pi}+\frac{4}{\pi}\theta(\sigma-|\Delta|)(\sigma-|\Delta|), \\ B &= \frac{\partial^2 V_3}{\partial\sigma\partial|\Delta|}=\frac{\partial^2 V_3}{\partial|\Delta|\partial\sigma}=\frac{4}{\pi}[\sigma-\theta(\sigma-|\Delta|)(\sigma-|\Delta|)], \\ C &= \frac{\partial^2 V_3}{\partial|\Delta|^2}=\frac{1}{2H_P}-\frac{4\Lambda_3}{\pi^2}+\frac{4|\Delta|}{\pi}+\frac{4}{\pi}\theta(\sigma-|\Delta|)(\sigma-|\Delta|). \end{aligned} \tag{20}$$

The equations (18) and (19) have the following four different solutions which will be discussed in order.

(i) $(\sigma,|\Delta|)$=(0,0). It is a maximum point of $V_3(\sigma,|\Delta|)$, since in this case we have

$$\begin{aligned} A &= 1/2G_S-2N_c\Lambda_3/\pi^2<0, \\ K &= A\left(1/2H_P-4\Lambda_3/\pi^2\right)>0, \end{aligned}$$

assuming Eqs. (18) and (19) have solutions of non-zero σ and $|\Delta|$.

(ii) $(\sigma,|\Delta|)$=(σ_1,0), where the non-zero σ_1 satisfies the equation

$$1/2G_S-2N_c\Lambda_3/\pi^2+N_c\sigma_1/\pi=0, \tag{21}$$

under the condition that $G_S\Lambda_3/\pi^2>1/4N_c$. For this solution, by means of Eq.(21) we obtain

$$\begin{aligned} A &= N_c\sigma_1/\pi, \\ K &= A\left(\frac{1}{2H_P}-\frac{1}{N_cG_S}+\frac{2\sigma_1}{\pi}\right)>0, \text{ if } \frac{G_S}{H_P}>\frac{2}{N_c}. \end{aligned}$$

Hence (σ_1,0) is a minimum point of $V_3(\sigma,|\Delta|)$ when $G_S/H_P>2/N_c$.

(iii) $(\sigma,\ |\Delta|)$= (0, Δ_1), where non-zero Δ_1 obeys the equation

$$1/2H_P-4\Lambda_3/\pi^2+2\Delta_1/\pi=0, \tag{22}$$

under the condition that $H_P\Lambda_3/\pi^2 > 1/8$. For this solution, by using Eq.(22), we get

$$\begin{aligned} A &= \frac{1}{2G_S} - \frac{N_c}{4H_P} + (4-N_c)\frac{\Delta_1}{\pi} && (23)\\ &= \frac{1}{2G_S} - \frac{2N_c\Lambda_3}{\pi^2} + \frac{4\Delta_1}{\pi}, && (24)\\ K &= 2\Delta_1 A/\pi. && (25) \end{aligned}$$

(a) When $N_c \leq 4$, from Eqs.(23) and (25) we have

$$A > 0, K > 0 \ \text{ for } \ \frac{G_S}{H_p} < \frac{2}{N_c}, \frac{H_P\Lambda_3}{\pi^2} > \frac{1}{8}.$$

(b) When $N_c > 4$, from Eqs.(24) and (25) we have

$$A > 0, K > 0 \ \text{ for } \ \frac{G_S\Lambda_3}{\pi^2} < \frac{1}{4N_c}, \frac{H_P\Lambda_3}{\pi^2} > \frac{1}{8}.$$

Noting that in this case the constraint $G_S/H_P < 2/N_c$ is also implied owing to that

$$1/8H_P < \Lambda_3/\pi^2 < 1/4N_cG_S.$$

Thus in the cases of both (a) and (b), the solution $(\sigma, |\Delta|) = (0, \Delta_1)$ is a minimum value point of $V_3(\sigma, |\Delta|)$.

(iv) $(\sigma, |\Delta|) = (\sigma_2, \Delta_2)$. In view of the function $\theta(\sigma - |\Delta|)$ in Eqs.(18) and (20), we have to consider the two cases of $\sigma_2 > \Delta_2$ and $\sigma_2 < \Delta_2$ respectively.

(a) $\sigma_2 > \Delta_2$. In this case Eqs.(18) and (19) will lead to

$$\frac{1}{2G_S} - \frac{N_c}{4H_P} = \frac{N_c\sigma_2^2 - 2\Delta_2^2}{\pi\sigma_2} > 0 \ \text{ thus } \ \frac{G_S}{H_P} < \frac{2}{N_c},$$

since $N_c \geq 2$. Then we obtain from Eq.(20) that

$$A = \frac{N_c\sigma_2^2 - 2\Delta_2^2}{\pi\sigma_2} > 0 \ \text{ and } \ K = -\frac{16\Delta_2^2}{\pi^2} < 0.$$

Hence the solution (σ_2, Δ_2) is not an extreme value point of $V_3(\sigma, |\Delta|)$ if $\sigma_2 > \Delta_2$.

(b) $\sigma_2 < \Delta_2$. In this case Eqs.(18) and (19) become

$$\frac{1}{2G_S} - \frac{2N_c\Lambda_3}{\pi^2} + \frac{4\Delta_2 + (N_c-2)\sigma_2}{\pi} = 0, \tag{26}$$

and

$$\frac{1}{2H_P} - \frac{4\Lambda_3}{\pi^2} + \frac{2\Delta_2}{\pi} + \frac{2\sigma_2^2}{\pi\Delta_2} = 0 \tag{27}$$

which may lead to the equations

$$\begin{aligned} \frac{1}{2H_P} - \frac{1}{N_cG_S} &= \frac{2}{\pi N_c\Delta_2}\left[(4-N_c)\Delta_2^2 + (N_c-2)\sigma_2\Delta_2 - N_c\sigma_2^2\right]\\ &= \frac{2(4-N_c)}{\pi N_c\Delta_2}\left[\left(\Delta_2 + \frac{N_c-2}{2(4-N_c)}\sigma_2\right)^2 + \frac{4N_c(N_c-4)-(N_c-2)^2}{4(4-N_c)^2}\sigma_2^2\right] && (28) \end{aligned}$$

and Eq.(20) will lead to that

$$\begin{aligned}K &= A\frac{2}{\pi\Delta_2}\left(\Delta_2^2-\sigma_2^2-\frac{8}{N_c-2}\Delta_2\sigma_2\right)\\ &= A\frac{2}{\pi\Delta_2}\left(\Delta_2+\frac{[16+(N_c-2)^2]^{\frac{1}{2}}-4}{N_c-2}\sigma_2\right)\tilde{K}\\ A &= (N_c-2)\frac{\sigma_2}{\pi}\\ \tilde{K} &= \Delta_2-\frac{[16+(N_c-2)^2]^{\frac{1}{2}}+4}{N_c-2}\sigma_2 \end{aligned}\tag{29}$$

Obviously, for $N_c > 2$, $\tilde{K} > 0$ will be the condition in which the solution (σ_2, Δ_2) becomes a minimum value point of $V_3(\sigma, |\Delta|)$. By means of Eqs.(28) and (29) we may discuss the relation between G_S and H_P when the (σ_2, Δ_2) becomes a possible minimum value point of $V_3(\sigma, |\Delta|)$. The result is obviously N_c-dependent.

(b1) $N_c = 2$. In this case $A = 0$, $K = -16\sigma_2^2/\pi^2 < 0$, so (σ_2, Δ_2) is not an extreme point of V_3.

(b2) $N_c = 3$. In this case, $\tilde{K} > 0$ or $K > 0$ become $\Delta_2 > (\sqrt{17}+4)\sigma_2$ or $\Delta_2^2 > \sigma_2^2 + 8\sigma_2\Delta_2$. Substituting these inequalities into Eq.(28) we get

$$\begin{aligned}\frac{1}{2H_P}-\frac{1}{N_cG_S} &= \frac{2}{3\pi\Delta_2}(\Delta_2^2+\sigma_2\Delta_2-3\sigma_2^2)\\ &> \frac{3\sigma_2^2}{\pi\Delta_2}(\sqrt{17}+4-\frac{2}{9}) > 0.\end{aligned}$$

(b3) $N_c = 4$. In this case, $\tilde{K} > 0$ becomes $\Delta_2 > (\sqrt{5}+2)\sigma_2$, then we have

$$\frac{1}{2H_P}-\frac{1}{N_cG_S} = \frac{\sigma_2}{\pi\Delta_2}(\Delta_2-2\sigma_2) > \frac{\sqrt{5}\sigma_2^2}{\pi\Delta_2} > 0.$$

The results in (b2) and (b3) indicate that when $N_c = 3, 4$ and $G_S/H_P > 2/N_c$, the non-zero solution (σ_2, Δ_2) of Eqs.(26) and (27) could be a minimum point of $V_3(\sigma, |\Delta|)$.

we have obtained in (ii) that when $G_S/H_P > 2/N_c$, the non-zero solution $(\sigma_1, 0)$ of Eqs.(18) and (19) is also a minimum point of $V_3(\sigma, |\Delta|)$ and the result is N_c-independent. In this case we must determine which one of the solutions (σ_2, Δ_2) and $(\sigma_1, 0)$ is the least value point of $V_3(\sigma, |\Delta|)$. For this purpose, we will compare the value of $V_3(\sigma, |\Delta|)$ at the point $(\sigma_1, 0)$ with the one at the point (σ_2, Δ_2). In fact, by using Eqs.(17) and (21), we may obtain that

$$V_3(\sigma_1, 0) = -N_c\sigma_1^3/6\pi \tag{30}$$

and by using Eqs.(17),(26)and (27), we can get that

$$V_3(\sigma_2, \Delta_2)|_{\sigma_2<\Delta_2} = -\frac{1}{3\pi}\left(\Delta_2^3+3\Delta_2\sigma_2^2+\frac{N_c-2}{2}\sigma_2^3\right). \tag{31}$$

In addition, comparing Eq.(21) with Eq.(26) with G_S and H_P being fixed, we will be led to the relation that

$$N_c\sigma_1 = 4\Delta_2 + (N_c - 2)\sigma_2 \tag{32}$$

By means of Eq.(32) we may calculate that

$$\begin{aligned} V_3(\sigma_1, 0) - V_3(\sigma_2, \Delta_2) &= -\frac{1}{3\pi}\begin{cases} \left[\frac{1}{9}(23\Delta_2^3 - 4\sigma_2^3) + \frac{1}{3}\sigma_2\Delta_2(8\Delta_2 - 7\sigma_2)\right], & \text{if } \ N_c = 3 \\ \left[\Delta_2^3 - \frac{3}{4}\sigma_2^3 + \frac{1}{2}\sigma_2\Delta_2(2\Delta_2 - \sigma_2)\right], & \text{if } \ N_c = 4 \end{cases} \\ &< 0 \quad \text{for } \ \sigma_2 < \Delta_2 \end{aligned} \tag{33}$$

This shows that when $N_c = 3, 4$ and $G_S/H_P > 2/N_c$, the least value point of $V_3(\sigma, |\Delta|)$ is $(\sigma_1, 0)$ rather than (σ_2, Δ_2). The same conclusion is also true for the case of $N_c = 2$ owing to the fact that (σ_2, Δ_2) is now not an extreme value point of $V_3(\sigma, |\Delta|)$, as shown in (b1). These results indicate that, when $G_S/H_P > 2/N_c$, only the pure quark-antiquark condensates could exist in the ground state of the 3D GN model.

(b4) $N_c > 4$. In this case, the solution (σ_2, Δ_2) which satisfies $\tilde{K} > 0$ will be a minimum value point of $V_3(\sigma, |\Delta|)$. Noting that the right-handed side of Eq.(28) is always less than zero when $N_c > 4$ and this implies that

$$G_S/H_P < 2/N_c \ \text{ when } \ N_c > 4,$$

hence we may affirm that if the solution (σ_2, Δ_2) is a minimum value point of $V_3(\sigma, |\Delta|)$ when $N_c > 4$, then the condition $G_S/H_P < 2/N_c$ must be obeyed.

Now the least value points of $V_3(\sigma, |\Delta|)$ can be summed up as follows:

$$(\sigma, |\Delta|) = \begin{cases} (0, \Delta_1), & \text{if } \ G_S/H_P < 2/N_c, \ H_P\Lambda_3/\pi^2 > 1/8, & & \text{for } N_c \le 4 \\ \begin{cases} (0, \Delta_1), \\ (\sigma_2, \Delta_2), \end{cases} & \text{if } \ G_S/H_P < 2/N_c, \ H_P\Lambda_3/\pi^2 > 1/8, & G_S\Lambda_3/\pi^2 \begin{cases} < 1/4N_c, \\ > 1/4N_c, \end{cases} & \text{for } N_c > 4 \\ (\sigma_1, 0), & \text{if } \ G_S/H_P > 2/N_c, & G_S\Lambda_3/\pi^2 > 1/4N_c, & \text{for all } N_c \end{cases} \tag{34}$$

Eq.(34) gives the $G_S - H_P$ phase diagrams Fig.3 and Fig.4 of the 3D GN model.

It can be seen from Eq.(34) and Figs.3-4 that, different from the 4D NJL and 2D GN model, the qualitative feature of fermionic condensates appearing in the ground state obviously depends on N_c. When the necessary conditions $G_S\Lambda_3 > \pi^2/4N_c$ and $H_P\Lambda_3 > \pi^2/8$ are satisfied, the ground state of the system could be in a pure quark-antiquark condensate phase only if $G_S/H_P > 2/N_c$ for all N_c, a similar result to the one obtained in the 4D and 2D model. However, if $G_S/H_P < 2/N_c$, then whether the coexistence phase with the condendates $\langle\bar{q}q\rangle$ and $\langle qq\rangle$ could appear will depend on N_c. The coexistence



Fig.3: G_S-H_P phase diagram of a 3D Gross-Neveu model when $N_c \leq 4$ with the denotations that $(0, \Delta_1)$–pure $\langle qq\rangle$ phase and $(\sigma_1, 0)$–pure $\langle\bar{q}q\rangle$ phase.



Fig.4: G_S-H_P phase diagram of a 3D Gross-Neveu model when $N_c \geq 5$ with the denotations that $(0, \Delta_1)$–pure $\langle qq\rangle$ phase, (σ_2, Δ_2)–mixed phase with $\langle\bar{q}q\rangle$ and $\langle qq\rangle$ and $(\sigma_1, 0)$–pure $\langle\bar{q}q\rangle$ phase.

phase could not exist if $N_c \leq 4$ and could do only if $N_c \geq 5$ in a 3D GN model. In addition, also similar to the 4D and 2D model, as N_c increases, the regions in the phase diagram occupied by the phases with the diquark condensates including the pure diquark condensate phase and the coexistence phase with quark-antiquark and diquark condensate will decrease gradually, and finally go to zeroes as $N_c \to \infty$.

V. CONCLUSIONS

We have researched interplay between the quark-antiquark and the diquark condensates in vacuum in two-flavor four-fermion interaction models with any color number N_c. The results are qualitatively consistent with the ones in $N_c = 3$ case in 4D NJL and 2D GN model, however different from the ones in $N_c = 3$ case in 3D GN model. It has been found that if the ratio of the four-fermion couplings of $(\bar{q}q)^2$-form and $(qq)^2$-form G_S/H_S (or G_S/H_P) $> 2/N_c$, the ratio of the color numbers of quarks entering the condensates $\langle qq\rangle$ and $\langle\bar{q}q\rangle$, (and also with sufficiently large G_S in 4D and 3D model), then only the pure $\langle\bar{q}q\rangle$ phase may exist and this conclusion is N_c-independent. Below $2/N_c$, (and also with sufficiently large H_S or H_P in 4D or 3D model), one will always first have a mixed phase with the condensates $\langle\bar{q}q\rangle$ and $\langle qq\rangle$, then a pure $\langle qq\rangle$ phase, except in the 3D GN model where the mixed phase does not appear if $N_c \leq 4$, the latter includes the known result in $N_c = 3$ case. A common characteristic in all the models is that as N_c increases, the phases with the diquark condensates $\langle qq\rangle$ including the pure $\langle qq\rangle$ phase and the mixed phase with the condensates $\langle\bar{q}q\rangle$ and $\langle qq\rangle$ will occupy smaller and smaller region in corresponding phase diagram and finally disappear as $N_c \to \infty$. In that limit only the quark-antiquark condensate phase become possible.

The above results certainly deepen our theoretical understanding of four-fermion interaction models. They not only may give a definite theoretical constraint on structure of some phenomenological models [12] but also could provide a useful clue to the feature of these models at finite temperature and finite quark chemical potential, as was shown in the 3D GN model with $N_c = 3$.

[1] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122** (1961) 345; **124** (1961) 246.

[2] D.J. Gross and A. Neveu, Phys. Rev. D **10** (1974) 3235.

[3] J. Goldstone, Nuovo Cimento **19** (1961) 154 .

[4] D. Weingarten, Phys. Rev. Lett. **51** (1983) 1830; S. Nussinov, Phys. Rev. Lett. **51** (1983) 2081; E. Witten, Phys. Rev. Lett. **51** (1983) 2351; C. Vafa and E. Witten, Nucl. Phys. B**234** (1984) 173.

[5] B. R. Zhou, Phys. Lett. B**215** (1988) 364; B**235** (1990) 412(E); Zhou Bang-Rong, Commun. Theor. Phys. (Beijing, China) **15** (1991) 319; **18** (1992) 347; **19** (1993) 377 .

[6] V. A. Miransky, *Dynamical symmetry breaking in Quantum Field Theory*, World Scientific, Singapore (1993) and the references therein.

[7] J. C. Collins and M. J. Perry, Phys. Rev. Lett. **34** (1975) 1353; M. Alford, K. Rajagopal, and F. Wilczek, Phys. Lett. B**422** (1998) 427; Nucl Phys. B**537** (1999) 443; R. Rapp, T. Schäfer, E. V. Shuryak, and M.Velkovsky, Phys. Rev. Lett. **81** (1998) 53; J. Berges and K. Rajagopal, Nucl. Phys. B**538** (1999) 215 .

[8] M. Buballa, Phys. Rep. **407** (2005) 205.

[9] D. Diakonov, H. Forkel, and M. Lutz, Phys. Lett.B **373** (1996) 147.

[10] B. Vanderheyden and A.D. Jackson, Phys. Rev. D**61** (2000) 076004.

[11] S. B. Rüster and D. H. Rischke, Phys. Rev. D**69** (2004) 045011.

[12] Zhou Bang-Rong, Commun. Theor. Phys. (Beijing, China) **47** (2007) 95.

[13] Zhou Bang-Rong, Commun. Theor. Phys. (Beijing, China) **47** (2007) 520.

[14] Zhou Bang-Rong, Commun. Theor. Phys. (Beijing, China) **47** (2007) 695.

[15] H. Kohyama, Phys. Rev. D**77** (2008) 045016.

[16] N. D. Mermin and H. Wagner, Phys. Rev. Lett. **17** (1966) 1133; S. Coleman, Commun. Math. Phys. **31** (1973) 259.